

Open Problems in Affine Algebraic Geometry,
in: *Affine Algebraic Geometry*,
J. Gutierrez, V. Shpilrain, J.-T. Yu, Eds.,
Contemp. Math., Amer. Math. Soc.
To appear.¹

PROBLEMS FOR PROBLEM SESSION

VLADIMIR L. POPOV

Below are the problems that I formulated at Open Problems Session of *Workshop on Group Actions on Rational Varieties*, McGill University and University of Montreal, Canada, March 2002.

Let k be an algebraically closed field of characteristic 0.

1. Roots of the Affine Cremona Group

Let $k^{[n]} := k[x_1, \dots, x_n]$ be the polynomial algebra in variables $x_1, \dots, x_n$ over k , and let $n > 1$. Put

$$D_i := \frac{\partial}{\partial x_i}.$$

The affine Cremona group $\text{Aut}_k k^{[n]}$ is an infinite dimensional algebraic group, and the set of ‘volume preserving’ transformations,

$$\text{Aut}_k^* k^{[n]} := \{\sigma \in \text{Aut}_k k^{[n]} \mid \det(D_i(\sigma(x_j)))_{1 \leq i, j \leq n} = 1\},$$

is its closed normal subgroup, [Sh], [Ka]. The group $\text{Aut}_k^* k^{[n]}$ is an infinite dimensional simple algebraic group, [Sh]. If D is a locally nilpotent k -derivation of $k^{[n]}$, then it is easily seen that $\exp tD \in \text{Aut}_k^* k^{[n]}$ for any $t \in k$, so D lies in $\text{Lie}(\text{Aut}_k^* k^{[n]})$.

It follows from [BB1], [BB2] that $n - 1$ is the maximum of dimensions of algebraic tori contained in $\text{Aut}_k^* k^{[n]}$, and that every algebraic torus in $\text{Aut}_k^* k^{[n]}$ of dimension $n - 1$ is conjugate to the ‘diagonal’ torus

$$T := \{\sigma := \text{diag}(t_1, \dots, t_n) \in \text{Aut}_k^* k^{[n]} \mid \sigma(x_i) = t_i x_i, t_i \in k, \prod_{i=1}^n t_i = 1\}.$$

Mimicing the notion of root from the theory of finite dimensional linear algebraic groups, cf., e.g., [Sp], introduce the following

¹Preprint: Open problems session of *Workshop on Group Actions on Rational Varieties*, McGill Univ. and Univ. of Montreal, Canada, March 2002 (G. Freudenburg, P. Russell, Eds.), Reports Centre Interuniv., Calcul. Math. Algèbr., Concordia, Laval, McGill, 2002–04, 13–14.

Definition 1.1. A nonzero locally nilpotent derivation D of $k^{[n]}$ is called a *root vector* of $\text{Aut}_k^* k^{[n]}$ with respect to T if there is a nontrivial character $\chi \in \text{Hom}(T, \mathbf{G}_m)$ such that

$$\sigma \circ D \circ \sigma^{-1} = \chi(\sigma)D \quad \forall \sigma \in T.$$

The character χ is called the *root* of $\text{Aut}_k^* k^{[n]}$ with respect to T corresponding to D .

Problem 1.1. Find all roots and root vectors of $\text{Aut}_k^* k^{[n]}$ with respect to T .

Among the root vectors there are evident ‘classical’ ones

$$x_i D_j, \text{ where } i \neq j, \text{ and } D_i$$

contained respectively in the Lie algebras of the special linear group SL_n (acting naturally on $k^{[n]}$ by linear transformations of $x_1, \dots, x_n$) and the group of translations

$$T_n := \{(x_1, \dots, x_n) \mapsto (x_1 + c_1, \dots, x_n + c_n) \mid c_i \in k\}.$$

The corresponding roots are respectively $\varepsilon_i \varepsilon_j^{-1}$ and ε_i where

$$\varepsilon_i(\text{diag}(t_1, \dots, t_n)) = t_i.$$

Problem 1.2. Describe the centralizer and the normalizer of T in $\text{Aut}_k^* k^{[n]}$.

Remark 1.1. The question whether $\text{Aut}_k^* k^{[n]}$ has no maximal (with respect to inclusion) algebraic tori of dimension $< n - 1$ is equivalent to asking whether every ‘volume preserving’ algebraic torus action on $\mathbf{A}^n$ is linearizable. The first nontrivial case is $n = 3$. Putting together the results of [KR], [KR1], [Ko], [KM], [KR2], [KR3] (see a historical account in [KMKR]) one obtains that in this case the answer is affirmative.

2. Generators of $\text{Aut}_k^* k^{[n]}$

Definition 2.1. A subgroup G of $\text{Aut}_k^* k^{[n]}$ is called *∂ -generated* if there exists a set $\mathcal{G}$ of locally nilpotent derivations of $k^{[n]}$ such that G is generated by the elements $\exp D$, where $D \in \mathcal{G}$.

Recall (cf. [P]) that, given a locally nilpotent derivation D of $k^{[n]}$, there is a simple method of constructing new such derivations: if f is an element of $\text{Ker } D := \{h \in k^{[n]} \mid Dh = 0\}$, then fD is a locally nilpotent derivation of $k^{[n]}$ as well. Thus for any such f we obtain an element $\exp fD$ of $\text{Aut}_k^* k^{[n]}$.

Definition 2.1. A subgroup G of $\text{Aut}_k^* k^{[n]}$ is called *finitely ∂ -generated* if there exists a finite set $\mathcal{G}$ of locally nilpotent derivations of $k^{[n]}$ such that G is generated by the elements $\exp fD$ where $D \in \mathcal{G}$ and $f \in \text{Ker } D$.

Example 2.1. Since the special linear group SL_n is generated by the root subgroups $\{\exp tx_i D_j \mid t \in k\}$, $1 \leq i \neq j \leq n$, cf. [Sp], it is finitely ∂ -generated (with $\mathcal{G} = \{D_1, \dots, D_n\}$). By the same reason, any connected semisimple algebraic subgroup G of $\text{Aut}_k^* k^{[n]}$ is finitely ∂ -generated. On the other hand, any algebraic torus in $\text{Aut}_k^* k^{[n]}$ is not ∂ -generated.

Example 2.2. Since the automorphism

$$(x_1, \dots, x_n) \mapsto (x_1 + c_1, \dots, x_n + c_n), \quad c_i \in k,$$

coincides with $(\exp c_1 D_1) \circ \dots \circ (\exp c_n D_n)$, the group of translations T_n is finitely ∂ -generated (with $\mathcal{G} = \{D_1, \dots, D_n\}$).

Example 2.3. Let $n > 1$. If $f_i \in k[x_{i+1}, \dots, x_n]$, $i = 1, \dots, n-1$, then $(\exp f_1 D_1) \circ \dots \circ (\exp f_{n-1} D_{n-1})$ is the automorphism

$$(x_1, \dots, x_n) \mapsto (x_1 + f_1, x_2 + f_2, \dots, x_{n-1} + f_{n-1}, x_n).$$

Hence the de Jonquière group J_n consisting of all such automorphisms is finitely ∂ -generated (with $\mathcal{G} = \{D_1, \dots, D_{n-1}\}$).

Problem 2.1. *Is the group $\text{Aut}_k^* k^{[n]}$ ∂ -generated? If yes, is it finitely ∂ -generated?*

Example 2.4. Since $\text{Aut}_k^* k^{[1]} = T_1$, it follows from Example 2.1 that the group $\text{Aut}_k^* k^{[1]}$ is finitely ∂ -generated (with $\mathcal{G} = \{D_1\}$).

Example 2.5. Since $\text{Aut}_k^* k^{[2]}$ is generated by SL_2 , T_2 and J_2 , cf. [vdK], it follows from Examples 2.1, 2.2 and 2.3 that the group $\text{Aut}_k^* k^{[2]}$ is finitely ∂ -generated (with $\mathcal{G} = \{D_1, D_2\}$).

Remark 2.1. Let $\sigma \in \text{Aut}_k^* k^{[3]}$ be the Nagata automorphism, [N],

$$(x_1, x_2, x_3) \mapsto (x_1 - 2x_2(x_3x_1 + x_2^2) - x_3(x_3x_1 + x_2^2)^2, x_2 + x_3(x_3x_1 + x_2^2), x_3).$$

By [SU1], [SU2], it is not tame, i.e., does not lie in the subgroup of $\text{Aut}_k^* k^{[3]}$ generated by SL_3 , T_3 and J_3 . However notice that $\sigma = \exp fD$, where D is the locally nilpotent derivation of $k^{[3]}$ given by $D := -2x_2D_1 + x_3D_2$, and $f := x_3x_1 + x_2^2 \in \text{Ker } D$.

3. Two Locally Nilpotent Derivations

Let P and Q be locally nilpotent derivations of $k^{[n]}$.

Problem 3.1. *When is the minimal closed subgroup of $\text{Aut}_k k^{[n]}$ containing the one-dimensional subgroups*

$$\{\exp tP \mid t \in k\} \text{ and } \{\exp tQ \mid t \in k\}$$

finite dimensional?

4. Rationality of Some Homogeneous Spaces

Let G be a connected semisimple algebraic group over k , and let $\sigma \in \text{Aut}_k G$ be an element of finite order. Consider the subgroup

$$G^\sigma := \{g \in G \mid \sigma(g) = g\}.$$

Conjecture 4.1. *The algebraic variety G/G^σ is rational.*

If $\text{ord } \sigma = 2$, then, by [V], the homogeneous space G/G^σ contains a dense open orbit of a Borel subgroup of G . Since any orbit of a connected solvable linear algebraic transformation group is rational, this shows that the conjecture is true for $\text{ord } \sigma = 2$.

REFERENCES

- [BB1] Białynicki–Birula, A., *Remarks on the action of an algebraic torus on k^n* , I, Bull. Acad. Polon. Sci. Ser. Sci. Math., Astr., Phys. **XIV** (1967), 177–188.
- [BB2] Białynicki–Birula, A., *Remarks on the action of an algebraic torus on k^n* , II, Bull. Acad. Polon. Sci. Ser. Sci. Math., Astr., Phys. **XV** (1967), 123–125.
- [Ka] Kambayashi, T., *Automorphism group of a polynomial ring and algebraic group action on an affine space*, J. of Algebra **60** (1979), 439–451.
- [KM] Kaliman, S., Makar–Limanov, L., *On the Russell–Koras contactible 3-folds*, J. Alg. Geom. **6** (1997), 247–268.
- [KMKR] Kaliman, S., Makar–Limanov, L., Koras, M., Russell, P., *$\mathbb{C}^*$ -actions on $\mathbb{C}^3$ are linearizable*, E. Res. Announc. **3** (1997), 63–71.
- [Ko] Koras, M., *A characterization of $\mathbb{C}^2/\mathbb{Z}_a$* , Comp. Math. **87** (1993), 241–267.
- [KR] Kambayashi, T., Russell, P., *On linearizing algebraic torus actions*, J. Pure Appl. Algebra **23** (1982), 243–250.
- [KR1] Koras, M., Russell, P., *$\mathbb{G}_m$ -actions on $\mathbf{A}^3$* , Can. Math. Soc. Conf. Proc. **10** (1989), 269–276.
- [KR2] Koras, M., Russell, P., *$\mathbb{C}^*$ -actions on $\mathbb{C}^3$: the smooth locus is not of hyperbolic type*, CICMA reports **06** (1996).
- [KR3] Koras, M., Russell, P., *Contractible 3-folds and $\mathbb{C}^*$ -actions on $\mathbb{C}^3$* , J. Alg. Geom. **6** (1997), 671–695.
- [N] Nagata, M., *On automorphism group of $k[x, y]$* , Lectures in Math. **5**, Dept. Math., Kyoto Univ., Kinokuniya, 1972.
- [P] Popov, V. L., *On actions of $\mathbf{G}_a$ on $\mathbf{A}^n$* , Lect. Notes in Math. **1271** (1987), Springer Verlag, 237–242.
- [Sh] Shafarevich, I. R., *On some infinite dimensional algebraic groups*, Rend. Math. e Appl. **25** (1966), no. 2, 208–212.
- [SU1] Shestakov, I. P., Umirbaev, U. U., *The Nagata automorphism is wild*, Proc. Natl. Acad. Sci. USA **100** (2003), no. 22, 12561–12563 (electronic).

- [SU2] Shestakov, I. P., Umirbaev, U. U., *The tame and the wild automorphisms of polynomial rings in three variables*, J. Amer. Math. Soc. **17** (2004), no. 1, 197–227.
- [Sp] Springer, T., *Linear Algebraic Groups*, Second Edition, Birkhäuser, 1998.
- [V] Vust, Th., *Opération de groupes réductifs dans un type de cônes presque homogènes*, Bull. Soc. Math. France **102** (1974), 317–334.
- [vdK] van der Kulk, W., *On polynomial rings in two variables*, Nieuw Arch. Wisk. **1** (1953), 33–41.

STEKLOV MATHEMATICAL INSTITUTE, RUSSIAN ACADEMY OF SCIENCES, GUBKINA 8,
MOSCOW, 117966, RUSSIA

E-mail address: `popovvl@orc.ru`